

Pion Production and the Pion Form Factor

H.P. Blok

Vrije Universiteit and NIKHEF, Amsterdam, The Netherlands

motivation

method

results and future

data on ^2H

high $|t|$

Hall C Workshop, August 20, 2004.

The Fpi2 Collaboration

D. Gaskell, M.K. Jones, D. Mack (Co-spokesman), J. Roche, G. Smith, W. Vulcan, G. Warren

Jefferson Lab, Newport News, VA, USA

E.J. Brash, G.M. Huber (Co-spokesman), V. Kovaltchouk, G.J. Lolos, C. Xu

University of Regina, Regina, SK, Canada

H. Blok (Co-spokesman), V. Tvaskis

Vrije Universiteit, Amsterdam, Netherlands

E. Beise, H. Breuer, C.C. Chang, T. Horn, P. King, J. Liu, P.G. Roos

University of Maryland, College Park, MD, USA

W. Boeglin, P. Markowitz, J. Reinhold

Florida International University, Miami, FL, USA

J. Arrington, R. Holt, D. Potterveld, P. Reimer, X. Zheng

Argonne National Laboratory, Argonne, IL, USA

H. Mkrtchyan, V. Tadevosyan

Yerevan Physics Institute, Yerevan, Armenia

S. Jin, W. Kim

Kyungook National University, Taegu, Korea

M.E. Christy, L.G. Tang

Hampton University, Hampton, VA, USA

J. Volmer

DESY, Hamburg, Germany

T. Miyoshi, Y. Okayasu, A. Matsumura

Tohoku University, Sendai, Japan

B. Barrett¹, A. Sarty¹, K. Aniol², D. Margaziotis², L. Pentchev³, C. Perdrisat³,

I. Niculescu⁴, V. Punjabi⁵, E. Gibson⁶

¹ *St. Mary's University, Halifax, NS, Canada* ² *California State University, Los Angeles, CA, USA*

³ *College of William and Mary, Williamsburg, VA, USA* ⁴ *George Washington University, Washington, DC, USA*


⁵ *Norfolk State University, Norfolk, VA, USA* ⁶ *California State University, Sacramento, CA, USA*

The Pion Form Factor in QCD

F_{π^+} has a unique place in our quest to understand hadronic structure, as the $\bar{q}q$ valence structure of the π^+ is relatively simple.

In quantum field theory, the pion form factor is given as the overlap integral

$$F_{\pi^+}(Q^2) = \int \phi_{\pi}^*(p) \phi_{\pi}(p+q) dp$$



The pion wave function can be separated into a ϕ_{π}^{soft} part with only low-momentum contributions ($k < k_0$) and a hard tail ϕ_{π}^{hard} . While ϕ_{π}^{hard} can be treated in pQCD, ϕ_{π}^{soft} cannot.

From a theoretical standpoint, the study of the Q^2 dependence of F_{π} focusses on finding a description for the soft and hard contributions to the pion wave function.


QCD Hard Scattering Picture:

At very high Q^2 , perturbative QCD (pQCD) can be used.

$$F_{\pi^+}(Q^2) = \int_0^1 dx \int_0^1 dy \frac{2g^2}{3xyQ^2} \phi(x)\phi(y)$$

$$g^2 = \frac{4}{3}\pi\alpha_s \quad (q - g \text{ coupling const})^2$$

$$xyQ^2 \quad \text{virtuality of exchanged gluon.}$$



As $Q^2 \rightarrow \infty$, only the hard portion of the wave function remains

$$\phi_\pi(x) \rightarrow 6f_\pi x(1-x)$$

where f_π determines the asymptotic normalization of the wave function from $\pi^+ \rightarrow \mu^+ \nu_\mu$ decay.



$$F_\pi \xrightarrow{Q^2 \rightarrow \infty} \frac{8\pi\alpha_s f_\pi^2}{Q^2}$$

This asymptotic normalization does **not** exist in the case of the nucleon form factors.

Intermediate Q^2 Scattering Picture:

At experimentally accessible Q^2 , the situation is more complicated

Both soft and hard components contribute.



[Bakulev, Passek-Kumericki, Schroers and Stefanis, hep-ph/0405062]

The soft contributions are poorly known, as is the interplay between the soft and hard contributions

[Braun, Khodjamirian, Maul, PRD **61**(00)073004.]


Determination of F_π via Pion Electroproduction

In the timelike region, F_π is determined from the $e^+e^- \rightarrow \pi^+\pi^-$ reaction. Our interest is in the spacelike region.


Up to $Q^2 = 0.3 \text{ GeV}^2$, F_π is measured directly from the scattering of 300 GeV pions from atomic electrons. These data determine the charge radius of the pion, $0.657 \pm 0.012 \text{ fm}$.

To access higher Q^2 , one must employ the $^1\text{H}(e, e'\pi^+)n$ reaction.

- At small $-t < 0.2 \text{ GeV}^2$, the t -channel diagram dominates σ_L .
- In the t -pole approximation, $\frac{d\sigma_L}{dt} \propto F_\pi^2$.
 - \Rightarrow In the actual extraction, a model incorporating the π^+ production mechanism and the effects of the ‘spectator’ nucleon is used to extract F_π from σ_L .
 - \Rightarrow π^+/π^- ratios from $^2\text{H}(e, e'\pi)$ are measured to test the validity of t -pole dominance and the model used.



What type of data do we need?



$$2\pi \frac{d^2\sigma}{dt d\phi} = \epsilon \frac{d\sigma_L}{dt} + \frac{d\sigma_T}{dt} + \sqrt{2\epsilon(\epsilon + 1)} \frac{d\sigma_{LT}}{dt} \cos\phi + \epsilon \frac{d\sigma_{TT}}{dt} \cos 2\phi$$

1. Take data at the smallest available $-t$, so that σ_L has maximum contribution from the π^+ pole.

\Rightarrow For a given Q^2 , higher W allows smaller $|t|_{min}$.

2. The extraction of F_π from σ_L requires that the $-t$ dependence of $d\sigma_L/dt$ is known.


\Rightarrow Only three of W , Q^2 , $-t$, and θ_π are independent.

Vary θ_π to obtain $-t$ dependence of the data.

\Rightarrow Since non-parallel data are needed, TT and LT must also be determined by the experiment.

Extraction of F_π from σ_L data

$Q^2 = 0.70 \text{ GeV}^2$ DESY expt [Z.Phys.C 3(79)101] used a Born term (BT) model, with modification to improve the description of the t -dependence of the data.



JLab expts use the Vanderhaeghen, Guidal and Laget Regge model, which provides a better treatment of the t -dependence.

“Model uncertainty” of extracted F_π result is obtained from fit to the t -dependence of the data under various assumptions.


DESY data also reanalyzed using Regge model:
 $\Rightarrow F_\pi(Q^2 = 0.70)$
increases by 0.05 from result obtained with BT model.

Recent and projected experimental data

E93-021 data is close to a monopole form factor with 0.657 fm charge radius.

These measurements were recently extended in a new Hall C experiment in the summer of 2003.

- $Q^2 = 2.45 \text{ GeV}^2$ using 6 GeV electron beam.
- Reduce model uncertainties in F_π extraction by obtaining data at higher $W = 2.21 \text{ GeV}$.
 - ⇒ New data will be closer to $t = m_\pi^2$ pole.
 - ⇒ Regge model can be applied with greater authority, so expect smaller model uncertainties.
- Q^2 region where F_π theoretical calculations begin to diverge.
 - ⇒ New data will constrain the treatment of soft contributions in QCD-based models.



Expect preliminary data to be released in second-half 2005.

F_π Long Term - 12 GeV Upgrade

The SHMS+HMS in Hall C will allow F_π to be measured to $Q^2 = 6 \text{ GeV}^2$, and possibly higher, depending on the favorability of the σ_L/σ_T ratio at large Q^2 .

The 5.5° forward angle capability of the SHMS is specifically driven by the requirement of the F_π experiment to access low $-t$.





Figure 4.3: Artist's Rendering of SHMS and HMS Spectrometers in Hall C. The top of the SOS Spectrometer is in the foreground.



These higher Q^2 data will have an unprecedented ability to test the state-of-the-art QCD calculations anticipated by that time.

Quasi-free pion production on ^2H .

Compare π^- and π^+ production

At low $|t|$: t-pole approximation: coupling to π^- or π^+

$$\Rightarrow R_{\pi^-/\pi^+} = 1.0$$

(isoscalar amplitudes are zero)

At high $|t|$: coupling to (charge of) quarks?




$$R_{\pi^-/\pi^+} = 1/4?$$

Q^2 dependence ?




L, T dependence ?

Preliminary results from Fpi1 (93-021) (Vardan Tadevosyan)

2001/09/11 08.56



DO NOT DISSEMINATE

E93021 R_T and VGL Regge


DO NOT DISSEMINATE

Pion production at high $|t|$.

Hall A results for photoproduction on ^2H .

Calculations by Huang, Jakob, Kroll and Passek-Kumericki

([hep-ph/0309071](#), Eur.Phys.J. C33 (2004) 91)



Suggests factorization into (non pQCD) photon-quark vertex and GPD.

$^1\text{H}(\text{e},\text{e}'\pi^+)\text{n}$ at high $|t|$.

Very preliminary results from a 'first look'.

Calculations by Jean-Marc Laget (hep-ph/0406153)

